Plaque Removal

in vivo study

Comparison of plaque removal in orthodontic subjects by Sonicare FlexCare and a manual toothbrush

DeLaurenti M, Putt M, Milleman J, Jenkins W, Wei J, Strate J. Plaque Removal by Sonicare and Manual Toothbrushes in Orthodontic Subjects. J Dent Res 87 (spec Iss B): 2044, 2008

Objective

To compare the plaque removal and gingivitis reduction ability for the Sonicare FlexCare (ProResults brush head) and Oral-B P-40[®] manual toothbrush in orthodontic population.

Methodology

Ninety-five healthy orthodontic subjects aged 12 years and older participated in a single-blind, randomized, parallel group clinical study assessing gingivitis and plaque over time for the Sonicare FlexCare and a manual toothbrush. All subjects were routine manual toothbrush users. At Visit 1, subjects were screened for eligibility (Modified Bonded Bracket Index (BBI) >2.0; Pocket depth ≤ 4mm). At Visit 2, subjects received a prophylaxis and were randomized to a treatment arm. Subjects were trained and given brushing instructions to use the assigned test device twice daily at home for two minutes. Subjects abstained from oral hygiene for 12-24 hours before each visit. At subsequent visits, safety and BBI were assessed before and after a two minute supervised brushing. Secondary efficacy measurements included Turesky-Modified Quigley-Hein Plaque Index (TPI) on tooth surfaces without brackets and a full-mouth Löe and Silness Gingival Index (LSGI). ANOVA was used for statistical analysis.

Results

The Sonicare FlexCare demonstrated superior plaque reduction in a single brushing on the buccal (bracketed) surfaces assessed by the Bonded Bracket Index compared to a manual toothbrush at the two-week and four-week evaluations (overall p<0.0001 at two weeks, p<0.0001 at four weeks).

The Sonicare FlexCare demonstrated significantly superior reduction in plaque over time (two and four weeks) on the lingual surfaces assessed by the Turesky-Modified Quigley-Hein Plaque Index compared to a manual toothbrush (overall p=0.0221 at two weeks and p=0.0025 at four weeks).

The Sonicare FlexCare demonstrated significantly superior plaque reduction in a single brushing on the lingual surfaces assessed by the Turesky-Modified Quigley-Hein Plaque Index compared to a manual toothbrush at the two-week and four-week evaluations (overall p=0.0001 at two weeks, overall p<0.0001 at four weeks).


The mean scoring Löe and Silness Gingival Index value was low (1.14, std 0.10) indicating that the population presented with healthy gingivae. Both groups, however, were able to demonstrate statistically significant improvement vs. baseline over time (overall p < 0.0001).

Conclusion

Sonicare FlexCare removed significantly more plaque than a manual toothbrush in a single brushing at two and four weeks on teeth with and without orthodontic brackets. Both toothbrushes were safe for use on oral soft tissues and orthodontic brackets.


Percent Plaque Reduction Bonded Bracket Index Single Brushing Bracketed Surfaces at 2 Weeks





Percent Plaque Reduction Single Brushing Bonded Bracket Index Bracketed Surfaces at 4 Weeks







Free Manuals Download Website

http://myh66.com

http://usermanuals.us

http://www.somanuals.com

http://www.4manuals.cc

http://www.manual-lib.com

http://www.404manual.com

http://www.luxmanual.com

http://aubethermostatmanual.com

Golf course search by state

http://golfingnear.com

Email search by domain

http://emailbydomain.com

Auto manuals search

http://auto.somanuals.com

TV manuals search

http://tv.somanuals.com